

Equivalent widths

- General methods
- Some examples

Equivalent widths are useful because can be easily converted into abundances, having a model atmosphere and the line parameters

Equivalent widths are independent of V_{sini} and V_{mac} broadening (you do not need to know V_{sini} and V_{mac} to measure abundances)


Equivalent widths

- General methods
 - **Direct integration**
 - **Line profile fitting**
- Some examples

Equivalent widths are useful because can be easily converted into abundances, having a model atmosphere and the line parameters

Equivalent widths are independent of V_{ini} and V_{mac} broadening (you do not need to know V_{ini} and V_{mac} to measure abundances)

Equivalent widths: definition and direct integration




Equivalent widths are measured in milliAngstroms

One directly measures the area between a spectral line and the continuum

Equivalent widths: line profile fitting

Spectral lines are typically Voigt profiles (convolution between Gaussian and Lorentzian profiles), however in most cases they can be approximated by either a gaussian or a rotational profile or a mix of the two




Binmag (IDL) by Oleg


<https://www.astro.uu.se/~oleg/binmag.html>

Visualisation, fitting of various parameters (e.g. V_r , V_{sini} , V_{mac}), measure equivalent widths using different methods, excellent for handling SB2, interface to SME for parameter and abundance determination


Equivalent widths: line profile fitting




Equivalent widths: line profile fitting




Equivalent widths: line profile fitting



Equivalent widths: line profile fitting



Equivalent widths: line profile fitting



Equivalent widths: line profile fitting

